



COMPOUNDING FORMULA

Diclofenac 10% in Diffusimax® 10 Kit Format: 100 g

CHEMICAL LIST

Drug	diclofenac	10.0 g
Wetting Agent	ethoxy diglycol	7.9 g (8.0 ml)
Base	Diffusimax10 Kit Compound A	15.2 g (16.1 ml)
	Diffusimax10 Kit Compound B	66.9 g

COMPOUNDING PROTOCOL

1.	Prepare Diffusimax10 cream. The amount of Diffusimax10 cream to make this formulation is 82.1 g, which requires 15.2 g (16.1 ml) Compound A and 66.9 g Compound B.
2.	In a mixing container, add 7.9 g (8.0 ml) ethoxy diglycol to 10.0 g diclofenac and mix to form a paste. There may be some granularity. This can be reduced by compressing the paste against the side of the mixing container using the mixing tool.
3.	Mix the paste intermittently over 5 minutes.
4.	q.s. the drug/wetting agent paste to 100 g with Diffusimax10 cream.
5.	Mix well.

DISPENSING INSTRUCTIONS

1.	Compounded formulation to be stored at room temperature. It may liquefy if cooled. Bringing it to room temperature and re-mixing will restore the cream.
----	--

BUD INFORMATION

Beyond use date: 180 days
Available at http://www.diffusimax.com/English/docs/BUD10_Diclofenac-10.pdf