



COMPOUNDING FORMULA

Lidocaine 10% in Diffusimax[®] 10 Kit Format: 100 g

CHEMICAL LIST

Drug	lidocaine	10.0 g
Wetting Agent	ethoxy diglycol	5.0 g (5.0 ml)
Base	Diffusimax10 Kit Compound A	15.7 g (16.7 ml)
	Diffusimax10 Kit Compound B	69.3 g

COMPOUNDING PROTOCOL

1.	Prepare Diffusimax10 cream. The amount of Diffusimax10 cream to make this formulation is 85.0 g, which requires 15.7 g (16.7 ml) Compound A and 69.3 g Compound B.
2.	In a mixing container, add 5.0 g (5.0 ml) ethoxy diglycol to 10.0 g lidocaine and mix to form a solution. There may be some granularity. This can be reduced by compressing any granules against the side of the mixing container using the mixing tool.
3.	q.s. the drug/wetting agent solution to 100 g with Diffusimax10 cream.
4.	Mix well.

DISPENSING INSTRUCTIONS

1.	Compounded formulation to be stored at room temperature. It may liquefy if cooled. Bringing it to room temperature and re-mixing will restore the cream.
----	--

BUD INFORMATION

Beyond use date: 180 days
Available at http://www.diffusimax.com/English/docs/BUD10_Lidocaine-10.pdf